



# NCERT Solutions for Class 10 Maths Unit 6

# **Triangles Class 10**

Unit 6 Triangles Exercise 6.1, 6.2, 6.3, 6.4, 6.5, 6.6 Solutions

Exercise 6.1: Solutions of Questions on Page Number: 122 Q1:

Fill in the blanks using correct word given in the brackets:-

(i) All circles are \_\_\_\_\_. (congruent, similar)

(ii) All squares are \_\_\_\_\_. (similar, congruent)

\_\_\_\_\_ triangles are similar. (isosceles, equilateral)

(iv) Two polygons of the same number of sides are similar, if (a) their corresponding angles are \_\_\_\_\_ and (b)

their corresponding sides are \_\_\_\_\_\_. (equal, proportional)

#### Answer:

- (i) Similar
- (ii) Similar
- (iii) Equilateral
- (iv) (a) Equal
- (b) Proportional


Q2:

Give two different examples of pair of

(i) Similar figures

(i)

Two



(ii)Non-similar figures


Answer:

Two equilateral triangles with sides 1 cm and 2 cm


squares with sides 1 cm and 2 cm








(ii) Trapezium and square




Triangle and parallelogram



### Q3:

State whether the following quadrilaterals are similar or not:



# Answer:


Quadrilateral PQRS and ABCD are not similar as their corresponding sides are proportional, i.e. 1:2, but their corresponding angles are not equal.

Exercise 6.2: Solutions of Questions on Page Number: 128 Q1:


In figure.6.17. (i) and (ii), DE || BC. Find EC in (i) and AD in (ii).

(i)






(ii)



# Answer:

(i)



Let EC = x cm

It is given that DE || BC.

By using basic proportionality theorem, we obtain

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\frac{1.5}{3} = \frac{1}{x}$$


$$x = \frac{3 \times 1}{1.5}$$

$$x = 2$$

(ii)







Let AD = x cm

It is given that DE || BC.

By using basic proportionality theorem, we obtain

$$\frac{AD}{DB} = \frac{AE}{EC}$$

$$\frac{x}{7.2} = \frac{1.8}{5.4}$$

$$x = \frac{1.8 \times 7.2}{5.4}$$


$$x = 2.4$$

$$\therefore AD = 2.4 \text{ cm}$$

#### Q2:

E and F are points on the sides PQ and PR respectively of a  $\Delta$ PQR. For each of the following cases, state whether EF || QR.

(i) PE = 
$$3.9$$
 cm, EQ =  $3$  cm, PF =  $3.6$  cm and FR =  $2.4$  cm



Given that, PE = 3.9 cm, EQ = 3 cm, PF = 3.6 cm, FR = 2.4 cm






$$\frac{PE}{EQ} = \frac{3.9}{3} = 1.3$$

$$\frac{PF}{FR} = \frac{3.6}{2.4} = 1.5$$

Hence, 
$$\frac{PE}{EQ} \neq \frac{PF}{FR}$$

Therefore, EF is not parallel to QR.




$$\frac{PE}{EQ} = \frac{4}{4.5} = \frac{8}{9}$$

$$\frac{PF}{FR} = \frac{8}{9}$$

Hence, 
$$\frac{PE}{EQ} = \frac{PF}{FR}$$

Therefore, EF is parallel to QR.





PE = 4 cm, QE = 4.5 cm, PF = 8 cm, RF = 9 cm



PQ = 1.28 cm, PR = 2.56 cm, PE = 0.18 cm, PF = 0.36 cm

$$\frac{PE}{PQ} = \frac{0.18}{1.28} = \frac{18}{128} = \frac{9}{64}$$

$$\frac{PF}{PR} = \frac{0.36}{2.56} = \frac{9}{64}$$


Hence, 
$$\frac{PE}{PQ} = \frac{PF}{PR}$$

Therefore, EF is parallel to QR.


# Q3:

In the following figure, if LM || CB and LN || CD, prove that

$$\frac{AM}{AB} = \frac{AN}{AD}$$



#### Answer:



In the given figure, LM || CB

By using basic proportionality theorem, we obtain





$$\frac{AM}{AB} = \frac{AL}{AC}$$

Similarly, LN || CD

$$\therefore \frac{AN}{AD} = \frac{AL}{AC}$$
 (ii)

From (i) and (ii), we obtain


$$\frac{AM}{AB} = \frac{AN}{AD}$$

# Q4:


In the following figure, DE || AC and DF || AE. Prove that

(i)

$$\frac{BF}{FE} = \frac{BE}{EC}$$



Answer:




$$\therefore \frac{BD}{DA} = \frac{BE}{EC}$$

(Basic Proportionality Theorem)

(*i*)







In  $\Delta BAE$ ,  $DF \parallel AE$ 

$$\therefore \frac{BD}{DA} = \frac{BF}{FE}$$

(Basic Proportionality Theorem)


(ii)

From (i) and (ii), we obtain


$$\frac{BE}{EC} = \frac{BF}{FE}$$

# Q5:

In the following figure, DE || OQ and DF || OR, show that EF || QR.



# Answer:




In  $\Delta$  POQ, DE || OQ

$$\therefore \frac{PE}{EO} = \frac{PD}{DO}$$

(Basic proportionality theorem) (i)







In ΔPOR, DF || OR


$$\therefore \frac{PF}{FR} = \frac{PD}{DO}$$

(Basic proportionality theorem) (ii)

From (i) and (ii), we obtain


$$\frac{PE}{EQ} = \frac{PF}{FR}$$

(Converse of basic proportionality theorem)



Q6:


In the following figure, A, B and C are points on OP, OQ and OR respectively such that AB || PQ and AC || PR. Show that BC || QR.



Answer:








In Δ POQ, AB || PQ

$$\therefore \frac{OA}{AP} = \frac{OB}{BQ}$$

(Basic proportionality theorem) (i)



In ΔPOR, AC || PR


$$\therefore \frac{OA}{AP} = \frac{OC}{CR}$$

(By basic proportionality theorem) (ii)

From (i) and (ii), we obtain

$$\frac{OB}{BQ} = \frac{OC}{CR}$$

(By the converse of basic proportionality theorem)




Q7:

Using Basic proportionality theorem, prove that a line drawn through the mid-points of one side of a triangle parallel to another side bisects the third side. (Recall that you have proved it in Class IX).



Answer:



Consider the given figure in which PQ is a line segment drawn through the mid-point P of line AB, such that  $PQ \parallel BC$ 

By using basic proportionality theorem, we obtain

$$\frac{AQ}{QC} = \frac{AP}{PB}$$

$$\frac{AQ}{QC} = \frac{1}{1}$$

(P is the mid-point of AB.  $\therefore$  AP = PB)


$$\Rightarrow$$
 AQ = QC

Or, Q is the mid-point of AC.

Q8:

Using Converse of basic proportionality theorem, prove that the line joining the mid-points of any two sides of a triangle is parallel to the third side. (Recall that you have done it in Class IX).

Answer:



Consider the given figure in which PQ is a line segment joining the mid-points P and Q of line AB and AC respectively.

i.e., AP = PB and AQ = QC

It can be observed that



$$\frac{AP}{PB} = \frac{1}{1}$$
and 
$$\frac{AQ}{QC} = \frac{1}{1}$$

$$\therefore \frac{AP}{PB} = \frac{AQ}{QC}$$

Hence, by using basic proportionality theorem, we obtain


PQ||BC

### Q9:

ABCD is a trapezium in which AB || DC and its diagonals intersect each other at the point O. Show

$$\frac{AO}{BO} = \frac{CO}{DO}$$
.

# Answer:



Draw a line EF through point O, such that  $EF \parallel CD$ 

By using basic proportionality theorem, we obtain

$$\frac{AE}{ED} = \frac{AO}{OC} \tag{1}$$

In 
$$\triangle ABD$$
, OE || AB

So, by using basic proportionality theorem, we obtain

$$\frac{ED}{AE} = \frac{OD}{BO}$$

$$\Rightarrow \frac{AE}{ED} = \frac{BO}{OD} \qquad (2)$$

From equations (1) and (2), we obtain





$$\frac{AO}{OC} = \frac{BO}{OD}$$

$$\Rightarrow \frac{AO}{BO} = \frac{OC}{OD}$$


Q10:

 $\frac{AO}{=}\frac{CO}{}$ 


The diagonals of a quadrilateral ABCD intersect each other at the point O such that  $\overline{BO}$   $\overline{DO}$  Show that ABCD is a trapezium.

#### Answer:

Let us consider the following figure for the given question.



Draw a line OE || AB



In ΔABD, OE || AB

By using basic proportionality theorem, we obtain

$$\frac{AE}{ED} = \frac{BO}{OD} \tag{1}$$

However, it is given that



$$\frac{AO}{OC} = \frac{OB}{OD}$$

(2)

From equations (1) and (2), we obtain

$$\frac{AE}{ED} = \frac{AO}{OC}$$


- ⇒ EO || DC [By the converse of basic proportionality theorem]
- ⇒ AB || OE || DC
- ⇒ AB || CD
- : ABCD is a trapezium.

Exercise 6.3: Solutions of Questions on Page Number: 138


Q1:


State which pairs of triangles in the following figure are similar? Write the similarity criterion used by you for answering the question and also write the pairs of similar triangles in the symbolic form:

(i)





(ii)






(iii)













(v)



(vi)



#### Answer:

(i) 
$$\angle A = \angle P = 60^{\circ}$$

$$\angle B = \angle Q = 80^{\circ}$$

$$\angle C = \angle R = 40^{\circ}$$

Therefore, ΔABC âˆÂ¼ ΔPQR [By AAA similarity criterion]

$$\frac{AB}{OR} = \frac{BC}{RP} = \frac{CA}{PO}$$

(ii)


- (iii)The given triangles are not similar as the corresponding sides are not proportional.
- (iv) In âˆâ€ MNL and âˆâ€ QPR, we observe that, MNQP = MLQR = 12

### Q2:

In the following figure,  $\triangle$ ODC  $\propto$ 1/4  $\triangle$ OBA,  $\angle$  BOC = 125° and  $\angle$  CDO = 70°. Find  $\angle$  DOC,  $\angle$  DCO and  $\angle$  OAB







### Answer:

DOB is a straight line.

$$\therefore$$
 ∠ DOC + ∠ COB = 180°

$$\Rightarrow$$
  $\angle$  DOC = 180 $^{\circ}$  - 125 $^{\circ}$ 

= 55°

In ΔDOC,

$$\angle$$
 DCO +  $\angle$  CDO +  $\angle$  DOC = 180°

(Sum of the measures of the angles of a triangle is 180°.)

$$\Rightarrow$$
  $\angle$  DCO + 70° + 55° = 180°

$$\Rightarrow$$
  $\angle$  DCO = 55°

It is given that  $\triangle$ ODC  $\angle$ Â $\frac{1}{4}$   $\triangle$ OBA.

 $\therefore$   $\angle$  OAB =  $\angle$  OCD [Corresponding angles are equal in similar triangles.]

$$\Rightarrow$$
  $\angle$  OAB = 55°


# Q3:

Diagonals AC and BD of a trapezium ABCD with AB || DC intersect each other at the point O. Using a

$$\frac{AO}{OC} = \frac{OB}{OD}$$

similarity criterion for two triangles, show that  $\overline{OC}$   $\overline{OD}$ 

# Answer:



In  $\Delta$ DOC and  $\Delta$ BOA,



∠CDO = ∠ABO [Alternate interior angles as AB || CD]

∠DCO = ∠BAO [Alternate interior angles as AB || CD]

∠DOC = ∠BOA [Vertically opposite angles]


 $\div$   $\Delta DOC~\tilde{A} \phi \ddot{E} \dagger \hat{A} \%~\Delta BOA~[AAA~similarity~criterion]$ 

$$\therefore \frac{DO}{BO} = \frac{OC}{OA}$$
$$\Rightarrow \frac{OA}{OC} = \frac{OB}{OD}$$


[Corresponding sides are proportional]

Q4:

 $\frac{QR}{QS} = \frac{QT}{PR} \ \ \text{and} \ \ \angle 1 = \angle 2.$  Show that  $\Delta PQS - \Delta TQR$ 



Answer:



In  $\triangle PQR$ ,  $\angle PQR = \angle PRQ$ 

∴ PQ = PR (i)

Given,



$$\frac{QR}{QS} = \frac{QT}{PR}$$

Using (i), we obtain

$$\frac{QR}{QS} = \frac{QT}{QP}$$

In  $\Delta PQS$  and  $\Delta TQR$ ,


$$\frac{QR}{OS} = \frac{QT}{OP}$$

$$\angle Q = \angle Q$$

# Q5:

S and T are point on sides PR and QR of  $\triangle$ PQR such that  $\angle$  P =  $\angle$  RTS. Show that  $\triangle$ RPQ  $\angle$ Â $\frac{1}{4}$   $\triangle$ RTS.

# Answer:



In ΔRPQ and ΔRST,

 $\angle$  RTS =  $\angle$  QPS (Given)

 $\angle R = \angle R$  (Common angle)


∴ ∆RPQ ∝¼ ∆RTS (By AA similarity criterion)

### Q6:

In the following figure, if  $\triangle ABE \cong \triangle ACD$ , show that  $\triangle ADE \propto \frac{1}{4} \triangle ABC$ .







# Answer:

It is given that  $\triangle ABE \cong \triangle ACD$ .

∴ AB = AC [By CPCT] (1)

And, AD = AE [By CPCT] (2)

In  $\triangle ADE$  and  $\triangle ABC$ ,


$$\frac{AD}{AB} = \frac{AE}{AC}$$
 [Dividing equation (2) by (1)]

 $\angle A = \angle A$  [Common angle]

∴ ΔADE âˆÂ¼ ΔABC [By SAS similarity criterion]

#### Q7:

In the following figure, altitudes AD and CE of  $\Delta ABC$  intersect each other at the point P. Show that:




- (i)  $\triangle$ AEP  $\propto \frac{1}{4} \triangle$ CDP
- (ii) ΔABD ∝¼ ΔCBE
- (iii) ΔAEP ∝¼ ΔADB
- (v)  $\triangle$ PDC  $\propto \frac{1}{4}$   $\triangle$ BEC

#### Answer:

(i)







In  $\triangle$ AEP and  $\triangle$ CDP,


∠ AEP = ∠ CDP (Each 90°)

∠ APE = ∠ CPD (Vertically opposite angles)

Hence, by using AA similarity criterion,

ΔAEP ∝¼ ΔCDP

(ii)



In ΔABD and ΔCBE,


∠ ADB = ∠ CEB (Each 90°)

 $\angle$  ABD =  $\angle$  CBE (Common)

Hence, by using AA similarity criterion,

ΔABD ∝¼ ΔCBE

(iii)



In  $\triangle AEP$  and  $\triangle ADB$ ,

 $\angle$  AEP =  $\angle$  ADB (Each 90°)

 $\angle$  PAE =  $\angle$  DAB (Common)


Hence, by using AA similarity criterion,

ΔAEP ∝¼ ΔADB

(iv)







In  $\triangle PDC$  and  $\triangle BEC$ ,

∠ PDC = ∠ BEC (Each 90°)

 $\angle$  PCD =  $\angle$  BCE (Common angle)


Hence, by using AA similarity criterion,

ΔPDC ∝¼ ΔBEC

#### Q8:

E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at F. Show that  $\triangle$ ABE  $\angle \hat{A}$  $^{\prime}_{A}$   $\triangle$ CFB

#### Answer:



In  $\triangle ABE$  and  $\triangle CFB$ ,

 $\angle A = \angle C$  (Opposite angles of a parallelogram)

∠ AEB = ∠ CBF (Alternate interior angles as AE || BC)


∴ ∆ABE ∝¼ ∆CFB (By AA similarity criterion)

### Q9:

In the following figure, ABC and AMP are two right triangles, right angled at B and M respectively, prove that:







(i) ΔABC âˆÂ¼ ΔAMP

$$\frac{CA}{PA} = \frac{BC}{MP}$$

### Answer:

In ΔABC and ΔAMP,

∠ABC = ∠AMP (Each 90°)

 $\angle A = \angle A$  (Common)

∴ ΔABC âˆÂ¼ ΔAMP (By AA similarity criterion)

$$\Rightarrow \frac{CA}{PA} = \frac{BC}{MF}$$

(Corresponding sides of similar triangles are proportional)


# Q10:

CD and GH are respectively the bisectors of  $\angle$ ACB and  $\angle$ EGF such that D and H lie on sides AB and FE of  $\triangle$ ABC and  $\triangle$ EFG respectively. If  $\triangle$ ABC  $\tilde{A}$ ¢ $\tilde{E}$ † $\hat{A}$ 1/4  $\triangle$ FEG, Show that:

$$\frac{CD}{GH} = \frac{AC}{FG}$$

- (ii) ΔDCB âˆÂ¼ ΔHGE
- (iii) ΔDCA âˆÂ¼ ΔHGF

# Answer:



It is given that ΔABC âˆÂ¼ ΔFEG.

$$\therefore \angle A = \angle F, \angle B = \angle E, \text{ and } \angle ACB = \angle FGE$$

∠ACB = ∠FGE

 $\therefore \angle ACD = \angle FGH$  (Angle bisector)

And,  $\angle DCB = \angle HGE$  (Angle bisector)

In  $\triangle$ ACD and  $\triangle$ FGH,

 $\angle A = \angle F$  (Proved above)

 $\angle$ ACD =  $\angle$ FGH (Proved above)

∴ ΔACD âˆÂ¼ ΔFGH (By AA similarity criterion)

$$\Rightarrow \frac{CD}{GH} = \frac{AC}{FG}$$

In ΔDCB and ΔHGE,

∠DCB = ∠HGE (Proved above)

 $\angle B = \angle E$  (Proved above)

∴ ΔDCB âˆÂ¼ ΔHGE (By AA similarity criterion)

In ΔDCA and ΔHGF,


 $\angle$ ACD =  $\angle$ FGH (Proved above)

 $\angle A = \angle F$  (Proved above)

∴ ΔDCA âˆÂ¼ ΔHGF (By AA similarity criterion)

# Q11:

In the following figure, E is a point on side CB produced of an isosceles triangle ABC with AB = AC. If AD  $\perp$  BC and EF  $\perp$  AC, prove that  $\Delta$ ABD  $\propto$ 1/4  $\Delta$ ECF



### Answer:

It is given that ABC is an isosceles triangle.

: AB = AC

 $\Rightarrow$   $\angle$  ABD =  $\angle$  ECF

In ΔABD and ΔECF,



∠ ADB = ∠ EFC (Each 90°)


 $\angle$  BAD =  $\angle$  CEF (Proved above)


∴ ΔABD ∠¼ ΔECF (By using AA similarity criterion)

### Q12:

Sides AB and BC and median AD of a triangle ABC are respectively proportional to sides PQ and QR and median PM of  $\Delta$ PQR (see the given figure). Show that  $\Delta$ ABC  $\angle$ Â $\frac{1}{4}$   $\Delta$ PQR.

#### Answer:





Median divides the opposite side.

$$BD = \frac{BC}{2}$$
 and  $QM = \frac{QR}{2}$ 

Given that,

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AD}{PM}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{\frac{1}{2}BC}{\frac{1}{2}QR} = \frac{AD}{PM}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM}$$

In  $\triangle ABD$  and  $\triangle PQM$ ,

$$\frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM}$$
 (Proved above)

∴ ΔABD âˆÂ¼ ΔPQM (By SSS similarity criterion)

⇒ ∠ABD = ∠PQM (Corresponding angles of similar triangles)

In  $\Delta ABC$  and  $\Delta PQR,$ 

 $\angle ABD = \angle PQM$  (Proved above)






$$\frac{AB}{PQ} = \frac{BC}{QR}$$

∴ ΔABC âˆÂ¼ ΔPQR (By SAS similarity criterion)

Q13:

D is a point on the side BC of a triangle ABC such that  $\angle ADC = \angle BAC$ . Show that  $CA^2 = CB.CD$ .

Answer:



In ΔADC and ΔBAC,

 $\angle ADC = \angle BAC$  (Given)

 $\angle ACD = \angle BCA$  (Common angle)

∴ ΔADC âˆÂ¼ ΔBAC (By AA similarity criterion)

We know that corresponding sides of similar triangles are in proportion.

$$\therefore \frac{CA}{CB} = \frac{CD}{CA}$$

$$\Rightarrow$$
 CA<sup>2</sup> = CB×CD


Q14:

Sides AB and AC and median AD of a triangle ABC are respectively proportional to sides PQ and PR and median PM of another triangle PQR. Show that  $\Delta ABC - \Delta PQR$ 

Answer:








Given that,

$$\frac{AB}{PQ} = \frac{AC}{PR} = \frac{AD}{PM}$$

Let us extend AD and PM up to point E and L respectively, such that AD = DE and PM = ML. Then, join B to E, C to E, Q to L, and R to L.



We know that medians divide opposite sides.

Therefore, BD = DC and QM = MR

Also, AD = DE (By construction)

And, PM = ML (By construction)

In quadrilateral ABEC, diagonals AE and BC bisect each other at point D.

Therefore, quadrilateral ABEC is a parallelogram.

∴ AC = BE and AB = EC (Opposite sides of a parallelogram are equal)

Similarly, we can prove that quadrilateral PQLR is a parallelogram and PR = QL, PQ = LR

It was given that

$$\frac{AB}{PQ} = \frac{AC}{PR} = \frac{AD}{PM}$$

$$AB \quad BE \quad 2$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BE}{QL} = \frac{2AD}{2PM}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BE}{QL} = \frac{AE}{PL}$$



∴ ΔABE âˆÂ¼ ΔPQL (By SSS similarity criterion)

We know that corresponding angles of similar triangles are equal.

$$\therefore \angle BAE = \angle QPL \dots (1)$$

Similarly, it can be proved that ΔAEC âˆÂ¼ ΔPLR and

$$\angle CAE = \angle RPL \dots (2)$$

Adding equation (1) and (2), we obtain

$$\angle BAE + \angle CAE = \angle QPL + \angle RPL$$

$$\Rightarrow \angle CAB = \angle RPQ \dots (3)$$

In  $\triangle$ ABC and  $\triangle$ PQR,

$$\frac{AB}{PQ} = \frac{AC}{PR}$$
 (Given)


 $\angle CAB = \angle RPQ$  [Using equation (3)]


∴ ΔABC âˆÂ¼ ΔPQR (By SAS similarity criterion)

### Q15:

A vertical pole of a length 6 m casts a shadow 4m long on the ground and at the same time a tower casts a shadow 28 m long. Find the height of the tower.

### Answer:





Let AB and CD be a tower and a pole respectively.

Let the shadow of BE and DF be the shadow of AB and CD respectively.

At the same time, the light rays from the sun will fall on the tower and the pole at the same angle.

Therefore,  $\angle DCF = \angle BAE$ 

And,  $\angle DFC = \angle BEA$ 

 $\angle$ CDF =  $\angle$ ABE (Tower and pole are vertical to the ground)

∴ ΔABE âˆÂ¼ ΔCDF (AAA similarity criterion)





$$\Rightarrow \frac{AB}{CD} = \frac{BE}{DF}$$

$$\Rightarrow \frac{AB}{6 \text{ m}} = \frac{28}{4}$$

$$\Rightarrow AB = 42 \text{ m}$$

Therefore, the height of the tower will be 42 metres.


#### Q16:

If AD and PM are medians of triangles ABC and PQR, respectively

$$\triangle ABC \sim \triangle PQR$$
 prove that  $t \frac{AB}{PQ} = \frac{AD}{PM}$ 

### Answer:

where



It is given that ΔABC âˆÂ¼ ΔPQR

We know that the corresponding sides of similar triangles are in proportion.

$$\frac{AB}{PQ} = \frac{AC}{PR} = \frac{BC}{QR} \dots (1)$$

Also, 
$$\angle A = \angle P$$
,  $\angle B = \angle Q$ ,  $\angle C = \angle R$  ... (2)

Since AD and PM are medians, they will divide their opposite sides.

BD = 
$$\frac{BC}{2}$$
 and QM =  $\frac{QR}{2}$  ... (3)

From equations (1) and (3), we obtain

$$\frac{AB}{PQ} = \frac{BD}{QM} \dots (4)$$

In  $\triangle ABD$  and  $\triangle PQM$ ,

 $\angle B = \angle Q$  [Using equation (2)]





$$\frac{AB}{PQ} = \frac{BD}{QM}$$
 [Using equation (4)]

∴ ΔABD âˆÂ¼ ΔPQM (By SAS similarity criterion)

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM}$$

Exercise 6.4: Solutions of Questions on Page Number: 143

Q1

Let  $\Delta ABC \sim \Delta DEF$  and their areas be, respectively, 64 cm² and 121 cm². If EF = 15.4 cm, find BC.

#### Answer:

It is given that  $\triangle ABC \sim \triangle DEF$ .

$$\therefore \frac{ar\left(\Delta ABC\right)}{ar\left(\Delta DEF\right)} = \left(\frac{AB}{DE}\right)^2 = \left(\frac{BC}{EF}\right)^2 = \left(\frac{AC}{DF}\right)^2$$

Given that,

$$EF = 15.4$$
 cm,

$$ar(\Delta ABC) = 64 \text{ cm}^2$$
,

$$ar(\Delta DEF) = 121 cm^2$$

$$\therefore \frac{\operatorname{ar}(ABC)}{\operatorname{ar}(DEF)} = \left(\frac{BC}{EF}\right)^{2}$$

$$\Rightarrow \left(\frac{64 \text{ cm}^{2}}{121 \text{ cm}^{2}}\right) = \frac{BC^{2}}{\left(15.4 \text{ cm}\right)^{2}}$$

$$\Rightarrow \frac{BC}{15.4} = \left(\frac{8}{11}\right) \text{cm}$$

$$\Rightarrow BC = \left(\frac{8 \times 15.4}{11}\right) \text{cm} = (8 \times 1.4) \text{cm} = 11.2 \text{ cm}$$


### Q2:

Diagonals of a trapezium ABCD with AB  $\parallel$  DC intersect each other at the point O. If AB = 2CD, find the ratio of the areas of triangles AOB and COD.

Answer:







Since AB || CD,

∴ ∠OAB = ∠OCD and ∠OBA = ∠ODC (Alternate interior angles)

In  $\triangle AOB$  and  $\triangle COD$ ,

 $\angle$ AOB =  $\angle$ COD (Vertically opposite angles)

∠OAB = ∠OCD (Alternate interior angles)

∠OBA = ∠ODC (Alternate interior angles)

∴ ΔAOB âˆÂ¼ ΔCOD (By AAA similarity criterion)

$$\therefore \frac{\operatorname{ar}(\Delta AOB)}{\operatorname{ar}(\Delta COD)} = \left(\frac{AB}{CD}\right)^2$$


Since AB = 2 CD,

$$\therefore \frac{\operatorname{ar}(\Delta AOB)}{\operatorname{ar}(\Delta COD)} = \left(\frac{2 \text{ CD}}{CD}\right)^2 = \frac{4}{1} = 4:1$$

# Q3:

In the following figure, ABC and DBC are two triangles on the same base BC. If AD intersects BC at O, show

$$\frac{\operatorname{area}(\Delta ABC)}{\operatorname{area}(\Delta DBC)} = \frac{AO}{DO}$$




### Answer:

Let us draw two perpendiculars AP and DM on line BC.







$$\frac{1}{2} \times Base \times Height$$
 We know that area of a triangle =

$$\therefore \frac{\operatorname{ar}(\Delta ABC)}{\operatorname{ar}(\Delta DBC)} = \frac{\frac{1}{2}BC \times AP}{\frac{1}{2}BC \times DM} = \frac{AP}{DM}$$

In ΔAPO and ΔDMO,

∠AOP = ∠DOM (Vertically opposite angles)

∴ ΔΑΡΟ Ã¢Ë†Â¼ ΔDMO (By AA similarity criterion)

$$\therefore \frac{AP}{DM} = \frac{AO}{DO}$$

$$\Rightarrow \frac{ar(\Delta ABC)}{ar(\Delta DBC)} = \frac{AO}{DO}$$

### Q4:

If the areas of two similar triangles are equal, prove that they are congruent.

# Answer:

Let us assume two similar triangles as ΔABC âˆÂ¼ ΔPQR.



$$\frac{\operatorname{ar}(\Delta ABC)}{\operatorname{ar}(\Delta PQR)} = \left(\frac{AB}{PQ}\right)^{2} = \left(\frac{BC}{QR}\right)^{2} = \left(\frac{AC}{PR}\right)^{2}$$
(1)

Given that, ar  $(\Delta ABC) = ar (\Delta PQR)$ 

$$\Rightarrow \frac{\operatorname{ar}(\Delta ABC)}{\operatorname{ar}(\Delta PQR)} = 1$$

Putting this value in equation (1), we obtain

$$1 = \left(\frac{AB}{PQ}\right)^2 = \left(\frac{BC}{QR}\right)^2 = \left(\frac{AC}{PR}\right)^2$$

 $\Rightarrow$  AB = PQ, BC = QR, and AC = PR


 $\triangle \Delta ABC \cong \Delta PQR$ 

(By SSS congruence criterion)

#### Q5:

D, E and F are respectively the mid-points of sides AB, BC and CA of  $\triangle$ ABC. Find the ratio of the area of  $\triangle$ DEF and  $\triangle$ ABC.

### Answer:



D and E are the mid-points of  $\triangle ABC$ .



$$\therefore$$
 DE || AC and DE =  $\frac{1}{2}$  AC

In ΔBED and ΔBCA,

$$\angle BED = \angle BCA$$
 (Corresponding angles)

$$\angle BDE = \angle BAC$$
 (Corresponding angles)

$$\angle EBD = \angle CBA$$
 (Common angles)

$$\frac{\operatorname{ar}(\Delta \operatorname{BED})}{\operatorname{ar}(\Delta \operatorname{BCA})} = \left(\frac{\operatorname{DE}}{\operatorname{AC}}\right)^2$$

$$\Rightarrow \frac{\operatorname{ar}(\Delta \operatorname{BED})}{\operatorname{ar}(\Delta \operatorname{BCA})} = \frac{1}{4}$$

$$\Rightarrow \operatorname{ar}(\Delta BED) = \frac{1}{4}\operatorname{ar}(\Delta BCA)$$

Similarly, 
$$ar(\Delta CFE) = \frac{1}{4}ar(CBA)$$
 and  $ar(\Delta ADF) = \frac{1}{4}ar(\Delta ABC)$ 

Also, 
$$ar(\Delta DEF) = ar(\Delta ABC) - [ar(\Delta BED) + ar(\Delta CFE) + ar(\Delta ADF)]$$


$$\Rightarrow \operatorname{ar}(\Delta DEF) = \operatorname{ar}(\Delta ABC) - \frac{3}{4}\operatorname{ar}(\Delta ABC) = \frac{1}{4}\operatorname{ar}(\Delta ABC)$$

$$\Rightarrow \frac{\text{ar}(\Delta \text{DEF})}{\text{ar}(\Delta \text{ABC})} = \frac{1}{4}$$

# Q6:

Prove that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding medians.

# Answer:



Let us assume two similar triangles as  $\triangle ABC \tilde{A} \phi \ddot{E} + \hat{A} \frac{1}{4} \triangle PQR$ . Let AD and PS be the medians of these triangles.

ΔABC âˆÂ¼ ΔPQR



$$\therefore \frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR}_{...(1)}$$

$$\angle A = \angle P$$
,  $\angle B = \angle Q$ ,  $\angle C = \angle R$  ... (2)

Since AD and PS are medians,

$$\therefore BD = DC = \frac{BC}{2}$$

And, 
$$QS = SR = 2$$

Equation (1) becomes

$$\frac{AB}{PQ} = \frac{BD}{QS} = \frac{AC}{PR}$$
... (3)

In  $\triangle$ ABD and  $\triangle$ PQS,

 $\angle B = \angle Q$  [Using equation (2)]

$$\frac{AB}{PO} = \frac{BD}{OS}$$

 $\frac{AB}{PQ} = \frac{BD}{QS}$  [Using equation (3)]

∴ ΔABD âˆÂ¼ ΔPQS (SAS similarity criterion)

Therefore, it can be said that

$$\frac{AB}{PQ} = \frac{BD}{QS} = \frac{AD}{PS} \dots (4)$$

$$\frac{\text{ar}\left(\Delta ABC\right)}{\text{ar}\left(\Delta PQR\right)} = \left(\frac{AB}{PQ}\right)^2 = \left(\frac{BC}{QR}\right)^2 = \left(\frac{AC}{PR}\right)^2$$

From equations (1) and (4), we may find that

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR} = \frac{AD}{PS}$$

And hence,


$$\frac{\operatorname{ar}(\Delta ABC)}{\operatorname{ar}(\Delta PQR)} = \left(\frac{AD}{PS}\right)^2$$

### Q7:

Prove that the area of an equilateral triangle described on one side of a square is equal to half the area of the equilateral triangle described on one of its diagonals.



Answer:



Let ABCD be a square of side a.

Therefore, its diagonal  $=\sqrt{2}a$ 

Two desired equilateral triangles are formed as  $\triangle ABE$  and  $\triangle DBF$ .

Side of an equilateral triangle,  $\triangle ABE$ , described on one of its sides = a

Side of an equilateral triangle,  $\Delta DBF$ , described on one of its diagonals  $=\sqrt{2}a$ 

We know that equilateral triangles have all its angles as 60 ° and all its sides of the same length. Therefore, all equilateral triangles are similar to each other. Hence, the ratio between the areas of these triangles will be equal to the square of the ratio between the sides of these triangles.

$$\frac{\text{Area of } \triangle \text{ ABE}}{\text{Area of } \triangle \text{DBF}} = \left(\frac{a}{\sqrt{2}a}\right)^2 = \frac{1}{2}$$

Q8:

ABC and BDE are two equilateral triangles such that D is the mid-point of BC. Ratio of the area of triangles ABC and BDE is

(A) 2:1

(B) 1:2


(C) 4:1

(D) 1:4

Answer:







We know that equilateral triangles have all its angles as 60 ° and all its sides of the same length. Therefore, all equilateral triangles are similar to each other. Hence, the ratio between the areas of these triangles will be equal to the square of the ratio between the sides of these triangles.

Let side of  $\triangle ABC = x$ 

$$\Delta BDE = \frac{x}{2}$$
 Therefore, side of

$$\therefore \frac{\text{area}(\Delta ABC)}{\text{area}(\Delta BDE)} = \left(\frac{x}{\frac{x}{2}}\right)^2 = \frac{4}{1}$$

Hence, the correct answer is (C).

Q9:

Sides of two similar triangles are in the ratio 4:9. Areas of these triangles are in the ratio

(A) 2:3

(B) 4:9

(C) 81:16

(D) 16:81

## Answer:

If two triangles are similar to each other, then the ratio of the areas of these triangles will be equal to the square of the ratio of the corresponding sides of these triangles.

It is given that the sides are in the ratio 4:9.

Therefore, ratio between areas of these triangles = 
$$\left(\frac{4}{9}\right)^2 = \frac{16}{81}$$

Hence, the correct answer is (D).

Exercise 6.5: Solutions of Questions on Page Number: 150

Q1:

Sides of triangles are given below. Determine which of them are right triangles? In case of a right triangle, write the length of its hypotenuse.

- (i) 7 cm, 24 cm, 25 cm
- (ii) 3 cm, 8 cm, 6 cm
- (iii) 50 cm, 80 cm, 100 cm
- (iv) 13 cm, 12 cm, 5 cm

#### Answer:

(i) It is given that the sides of the triangle are 7 cm, 24 cm, and 25 cm.

Squaring the lengths of these sides, we will obtain 49, 576, and 625.

Or. 
$$7^2 + 24^2 = 25^2$$

The sides of the given triangle are satisfying Pythagoras theorem.

Therefore, it is a right triangle.

We know that the longest side of a right triangle is the hypotenuse.

Therefore, the length of the hypotenuse of this triangle is 25 cm.

(ii) It is given that the sides of the triangle are 3 cm, 8 cm, and 6 cm.

Squaring the lengths of these sides, we will obtain 9, 64, and 36.

However, 9 + 36 ≠ 64

Or, 
$$3^2 + 6^2 \neq 8^2$$

Clearly, the sum of the squares of the lengths of two sides is not equal to the square of the length of the third side.

Therefore, the given triangle is not satisfying Pythagoras theorem.

Hence, it is not a right triangle.

(iii)Given that sides are 50 cm, 80 cm, and 100 cm.

Squaring the lengths of these sides, we will obtain 2500, 6400, and 10000.

However,  $2500 + 6400 \neq 10000$ 

Or, 
$$50^2 + 80^2 \neq 100^2$$

Clearly, the sum of the squares of the lengths of two sides is not equal to the square of the length of the third side.

Therefore, the given triangle is not satisfying Pythagoras theorem.

Hence, it is not a right triangle.

(iv)Given that sides are 13 cm, 12 cm, and 5 cm.



Squaring the lengths of these sides, we will obtain 169, 144, and 25.

Clearly, 144 + 25 = 169

Or, 
$$12^2 + 5^2 = 13^2$$

The sides of the given triangle are satisfying Pythagoras theorem.

Therefore, it is a right triangle.


We know that the longest side of a right triangle is the hypotenuse.

Therefore, the length of the hypotenuse of this triangle is 13 cm.

## Q2:

PQR is a triangle right angled at P and M is a point on QR such that PM  $\perp$  QR. Show that PM<sup>2</sup> = QM x MR.

## Answer:





Let 
$$\angle MPR = x$$

In AMPR,

$$\angle MRP = 180^{\circ} - 90^{\circ} - x$$

$$\angle MRP = 90^{\circ} - x$$

Similarly, in AMPQ,

$$\angle$$
MPQ = 90° - $\angle$ MPR

$$=90^{\circ}-x$$

$$\angle MQP = 180^{\circ} - 90^{\circ} - (90^{\circ} - x)$$

$$\angle MQP = x$$

In  $\triangle$  QMP and  $\triangle$ PMR,

$$\angle MPQ = \angle MRP$$

$$\angle PMQ = \angle RMP$$

$$\angle MQP = \angle MPR$$


$$\Rightarrow \frac{QM}{PM} = \frac{MP}{MR}$$

$$\Rightarrow PM^2 = QM \times MR$$

## Q3:

ABC is an isosceles triangle right angled at C. prove that  $AB^2 = 2 AC^2$ .

# Answer:



Given that  $\triangle ABC$  is an isosceles triangle.


Applying Pythagoras theorem in  $\Delta ABC$  (i.e., right-angled at point C), we obtain

$$AC^{2} + CB^{2} = AB^{2}$$
  
 $\Rightarrow AC^{2} + AC^{2} = AB^{2}$  (AC=CB)  
 $\Rightarrow 2AC^{2} = AB^{2}$ 

## Q4:

ABC is an isosceles triangle with AC = BC. If  $AB^2 = 2 AC^2$ , prove that ABC is a right triangle.

# Answer:



Given that,

$$AB^{2} = 2AC^{2}$$

$$\Rightarrow AB^{2} = AC^{2} + AC^{2}$$

$$\Rightarrow AB^{2} = AC^{2} + BC^{2} \text{ (As AC = BC)}$$


The triangle is satisfying the pythagoras theorem.

Therefore, the given triangle is a right - angled triangle.

# Q5:

ABC is an equilateral triangle of side 2a. Find each of its altitudes.

### Answer:



Let AD be the altitude in the given equilateral triangle,  $\Delta ABC$ .

We know that altitude bisects the opposite side.

$$\therefore$$
 BD = DC =  $a$ 



In ΔADB,

$$\angle ADB = 90^{\circ}$$

Applying pythagoras theorem, we obtain

$$AD^2 + DB^2 = AB^2$$

$$\Rightarrow AD^2 + a^2 = (2a)^2$$

$$\Rightarrow$$
 AD<sup>2</sup> +  $a^2$  =  $4a^2$ 

$$\Rightarrow AD^2 = 3a^2$$

$$\Rightarrow$$
 AD =  $a\sqrt{3}$ 


In an equilateral triangle, all the altitudes are equal in length.

Therefore, the length of each altitude will be  $\sqrt{3}a$ .

# Q6:

Prove that the sum of the squares of the sides of rhombus is equal to the sum of the squares of its diagonals.

## Answer:



In ΔΑΟΒ, ΔΒΟC, ΔCOD, ΔΑΟD,

Applying Pythagoras theorem, we obtain



$$AB^2 = AO^2 + OB^2$$
 ... (1)

$$BC^2 = BO^2 + OC^2$$
 ... (2)

$$CD^2 = CO^2 + OD^2$$
 ... (3)

$$AD^2 = AO^2 + OD^2$$
 ... (4)

Adding all these equations, we obtain

$$AB^{2} + BC^{2} + CD^{2} + AD^{2} = 2(AO^{2} + OB^{2} + OC^{2} + OD^{2})$$

$$=2\left[\left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2 + \left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2\right]$$


(Diagonals bisect each other)

$$=2\left(\frac{\left(AC\right)^2}{2} + \frac{\left(BD\right)^2}{2}\right)$$

$$= (AC)^2 + (BD)^2$$

# Q7:

In the following figure, O is a point in the interior of a triangle ABC, OD  $\perp$  BC, OE  $\perp$  AC and OF  $\perp$  AB. Show that



(i) 
$$OA^2 + OB^2 + OC^2 - OD^2 - OE^2 - OF^2 = AF^2 + BD^2 + CE^2$$


(ii) 
$$AF^2 + BD^2 + CE^2 = AE^2 + CD^2 + BF^2$$

# Answer:

Join OA, OB, and OC.







(i) Applying Pythagoras theorem in  $\triangle AOF$ , we obtain

$$OA^2 = OF^2 + AF^2$$

Similarly, in ΔBOD,

$$OB^2 = OD^2 + BD^2$$

Similarly, in ΔCOE,

$$OC^2 = OE^2 + EC^2$$

Adding these equations,

$$OA^{2} + OB^{2} + OC^{2} = OF^{2} + AF^{2} + OD^{2} + BD^{2} + OE^{2} + EC^{2}$$

$$OA^{2} + OB^{2} + OC^{2} - OD^{2} - OE^{2} - OF^{2} = AF^{2} + BD^{2} + EC^{2}$$

(ii) From the above result,


$$AF^{2} + BD^{2} + EC^{2} = (OA^{2} - OE^{2}) + (OC^{2} - OD^{2}) + (OB^{2} - OF^{2})$$

$$AF^2 + BD^2 + EC^2 = AE^2 + CD^2 + BF^2$$

# Q8:

A ladder 10 m long reaches a window 8 m above the ground. Find the distance of the foot of the ladder from base of the wall.

# Answer:



Let OA be the wall and AB be the ladder.



Therefore, by Pythagoras theorem,

$$AB^{2} = OA^{2} + BO^{2}$$

$$(10 \text{ m})^{2} = (8 \text{ m})^{2} + OB^{2}$$

$$100 \text{ m}^{2} = 64 \text{ m}^{2} + OB^{2}$$

$$OB^{2} = 36 \text{ m}^{2}$$

$$OB = 6 \text{ m}$$


Therefore, the distance of the foot of the ladder from the base of the wall is

6 m.

#### Q9:

A guy wire attached to a vertical pole of height 18 m is 24 m long and has a stake attached to the other end. How far from the base of the pole should the stake be driven so that the wire will be taut?

## Answer:



Let OB be the pole and AB be the wire.

By Pythagoras theorem,

$$AB^{2} = OB^{2} + OA^{2}$$

$$(24 \text{ m})^{2} = (18 \text{ m})^{2} + OA^{2}$$

$$OA^{2} = (576 - 324) \text{ m}^{2} = 252 \text{ m}^{2}$$

$$OA = \sqrt{252} \text{ m} = \sqrt{6 \times 6 \times 7} \text{ m} = 6\sqrt{7} \text{ m}$$

Therefore, the distance from the base is  $6\sqrt{7}$  m.

## Q10:


An aeroplane leaves an airport and flies due north at a speed of 1,000 km per hour. At the same time, another aeroplane leaves the same airport and flies due west at a speed of 1,200 km per hour. How far apart will be

the two planes after  $1\frac{1}{2}$  hours'





Answer:



Distance travelled by the plane flying towards north in  $1\frac{1}{2}hrs = 1,000 \times 1\frac{1}{2} = 1,500 \text{ km}$ 

Similarly, distance travelled by the plane flying towards west in  $1\frac{1}{2}$  hrs  $=1,200 \times 1\frac{1}{2}$  = 1,800 km

Let these distances be represented by OA and OB respectively.

Applying Pythagoras theorem,

Distance between these planes after  $\frac{1\frac{1}{2} \text{ hrs}}{\text{AB}} = \sqrt{\text{OA}^2 + \text{OB}^2}$   $= \left(\sqrt{(1,500)^2 + (1,800)^2}\right) \text{km} = \left(\sqrt{2250000 + 3240000}\right) \text{km}$   $= \left(\sqrt{5490000}\right) \text{km} = \left(\sqrt{9 \times 610000}\right) \text{km} = 300\sqrt{61} \text{ km}$ 

Therefore, the distance between these planes will be  $300\sqrt{61}$  km after  $1\frac{1}{2}$  hrs


# Q11:

Two poles of heights 6 m and 11 m stand on a plane ground. If the distance between the feet of the poles is 12 m, find the distance between their tops.

Answer:







Let CD and AB be the poles of height 11 m and 6 m.

Therefore, CP = 11 - 6 = 5 m

From the figure, it can be observed that AP = 12m

Applying Pythagoras theorem for ΔAPC, we obtain


$$AP^{2} + PC^{2} = AC^{2}$$
  
 $(12 \text{ m})^{2} + (5 \text{ m})^{2} = AC^{2}$   
 $AC^{2} = (144 + 25) \text{ m}^{2} = 169 \text{ m}^{2}$   
 $AC = 13 \text{ m}$ 

Therefore, the distance between their tops is 13 m.

# Q12:

D and E are points on the sides CA and CB respectively of a triangle ABC right angled at C. Prove that  $AE^2+BD^2=AB^2+DE^2$ 

## Answer:



Applying Pythagoras theorem in ΔACE, we obtain



$$AC^2 + CE^2 = AE^2$$
 ... (1)

Applying Pythagoras theorem in ΔBCD, we obtain

$$BC^2 + CD^2 = BD^2$$
 ... (2)

Using equation (1) and equation (2), we obtain

$$AC^2 + CE^2 + BC^2 + CD^2 = AE^2 + BD^2$$
 ... (3)

Applying Pythagoras theorem in  $\Delta CDE$ , we obtain

$$DE^2 = CD^2 + CE^2$$

Applying Pythagoras theorem in ΔABC, we obtain


$$AB^2 = AC^2 + CB^2$$

Putting the values in equation (3), we obtain

$$DE^2 + AB^2 = AE^2 + BD^2$$

### Q13:

The perpendicular from A on side BC of a  $\triangle$ ABC intersect BC at D such that DB = 3 CD. Prove that 2 AB<sup>2</sup> = 2 AC<sup>2</sup> + BC<sup>2</sup>



# Answer:

Applying Pythagoras theorem for ΔACD, we obtain

$$AC^2 = AD^2 + DC^2$$

$$AD^2 = AC^2 - DC^2$$
 ... (1)

Applying Pythagoras theorem in  $\triangle ABD$ , we obtain



$$AB^2 = AD^2 + DB^2$$

$$AD^2 = AB^2 - DB^2$$

From equation (1) and equation (2), we obtain

$$AC^2 - DC^2 = AB^2 - DB^2$$
 ... (3

It is given that 3DC = DB

$$\therefore DC = \frac{BC}{4} \text{ and } DB = \frac{3BC}{4}$$

Putting these values in equation (3), we obtain

$$AC^2 - \left(\frac{BC}{4}\right)^2 = AB^2 - \left(\frac{3BC}{4}\right)^2$$

$$AC^2 - \frac{BC^2}{16} = AB^2 - \frac{9BC^2}{16}$$

$$16AC^2 - BC^2 = 16AB^2 - 9BC^2$$


$$16AB^2 - 16AC^2 = 8BC^2$$

$$2AB^2 = 2AC^2 + BC^2$$

Q14:

In an equilateral triangle ABC, D is a point on side BC such that BD =  $\frac{1}{3}$  BC. Prove that 9 AD<sup>2</sup> = 7 AB<sup>2</sup>.

Answer:



Let the side of the equilateral triangle be a, and AE be the altitude of  $\triangle$ ABC.

$$\therefore BE = EC = \frac{BC}{2} = \frac{a}{2}$$

And, AE = 
$$\frac{a\sqrt{3}}{2}$$

Given that, BD = 
$$\frac{1}{3}$$
 BC





$$\therefore BD = \frac{a}{3}$$

$$\frac{a}{2} - \frac{a}{3} = \frac{a}{6}$$

Applying Pythagoras theorem in  $\Delta ADE$ , we obtain

$$AD^2 = AE^2 + DE^2$$


$$AD^{2} = \left(\frac{a\sqrt{3}}{2}\right)^{2} + \left(\frac{a}{6}\right)^{2}$$
$$= \left(\frac{3a^{2}}{4}\right) + \left(\frac{a^{2}}{36}\right)$$
$$= \frac{28a^{2}}{36}$$
$$= \frac{7}{9}AB^{2}$$

$$\Rightarrow$$
 9 AD<sup>2</sup> = 7 AB<sup>2</sup>

## Q15:

In an equilateral triangle, prove that three times the square of one side is equal to four times the square of one of its altitudes.

# Answer:



Let the side of the equilateral triangle be a, and AE be the altitude of  $\Delta ABC$ .

$$\therefore BE = EC = \frac{BC}{2} = \frac{a}{2}$$

Applying Pythagoras theorem in  $\Delta ABE$ , we obtain

$$AB^2 = AE^2 + BE^2$$



$$a^2 = AE^2 + \left(\frac{a}{2}\right)^2$$

$$AE^2 = a^2 - \frac{a^2}{4}$$

$$AE^2 = \frac{3a^2}{4}$$

$$4AE^{2} = 3a^{2}$$

 $\Rightarrow$  4 × (Square of altitude) = 3 × (Square of one side)

### Q16:


Tick the correct answer and justify: In  $\triangle ABC$ ,  $AB = 6\sqrt{3}$  cm, AC = 12 cm and BC = 6 cm.

The angle B is:

(A) 120° (B) 60°

(C) 90° (D) 45°

### Answer:



Given that, AB = 
$$6\sqrt{3}$$
 cm, AC = 12 cm, and BC = 6 cm

It can be observed that

$$AB^2 = 108$$

$$AC^2 = 144$$

And, 
$$BC^2 = 36$$

$$AB^2 + BC^2 = AC^2$$

The given triangle,  $\Delta ABC,$  is satisfying Pythagoras theorem.

Therefore, the triangle is a right triangle, right-angled at B.

$$∴$$
 ∠B = 90°

Hence, the correct answer is (C).




Exercise 6.6: Solutions of Questions on Page Number: 152


Q1:

$$\frac{QS}{SR} = \frac{PQ}{PR}$$

In the given figure, PS is the bisector of  $\angle QPR$  of  $\Delta PQR$ . Prove that  $\overline{SR} = \overline{PR}$ .



# Answer:



Let us draw a line segment RT parallel to SP which intersects extended line segment QP at point T.

Given that, PS is the angle bisector of ∠QPR.

$$\angle QPS = \angle SPR \dots (1)$$

By construction,

$$\angle$$
SPR =  $\angle$ PRT (As PS || TR) ... (2)

$$\angle QPS = \angle QTR \text{ (As PS || TR) ... (3)}$$

Using these equations, we obtain

$$\angle PRT = \angle QTR$$

By construction,

PS || TR






By using basic proportionality theorem for  $\Delta \text{QTR}, \\ \text{QSSR=QPPT}$ 

⇒QSSR

### Q2:


In the given figure, D is a point on hypotenuse AC of  $\triangle$ ABC, DM  $\bot$  BC and DN  $\bot$  AB, Prove that:

- (i)  $DM^2 = DN.MC$
- (ii)  $DN^2 = DM.AN$



### Answer:

(i)Let us join DB.



We have, DN || CB, DM || AB, and  $\angle$ B = 90°

- : DMBN is a rectangle.
- ∴ DN = MB and DM = NB

The condition to be proved is the case when D is the foot of the perpendicular drawn from B to AC.

$$\Rightarrow$$
  $\angle$ 2 +  $\angle$ 3 = 90° ... (1)

In ΔCDM,

$$\angle 1 + \angle 2 + \angle DMC = 180^{\circ}$$

$$\Rightarrow \angle 1 + \angle 2 = 90^{\circ} \dots (2)$$



In ΔDMB,

$$\angle 3 + \angle DMB + \angle 4 = 180^{\circ}$$

$$\Rightarrow \angle 3 + \angle 4 = 90^{\circ} \dots (3)$$

From equation (1) and (2), we obtain

$$\angle 1 = \angle 3$$

From equation (1) and (3), we obtain

In ΔDCM and ΔBDM,

 $\angle 1 = \angle 3$  (Proved above)

 $\angle 2 = \angle 4$  (Proved above)

∴ ΔDCM âˆÂ¼ ΔBDM (AA similarity criterion)

$$\Rightarrow \frac{BM}{DM} = \frac{DM}{MC}$$

$$\Rightarrow \frac{DN}{DM} = \frac{DM}{MC}$$
(BM = DN)

$$\Rightarrow$$
 DM<sup>2</sup> = DN  $\times$  MC

(ii) In right triangle DBN,

$$\angle 5 + \angle 7 = 90^{\circ} \dots (4)$$

In right triangle DAN,

$$\angle 6 + \angle 8 = 90^{\circ} \dots (5)$$

D is the foot of the perpendicular drawn from B to AC.

$$\Rightarrow \angle 5 + \angle 6 = 90^{\circ} \dots (6)$$

From equation (4) and (6), we obtain

From equation (5) and (6), we obtain

$$\angle 8 = \angle 5$$

In  $\triangle$ DNA and  $\triangle$ BND,

$$\angle 6 = \angle 7$$
 (Proved above)

$$\angle 8 = \angle 5$$
 (Proved above)

∴ ΔDNA âˆÂ¼ ΔBND (AA similarity criterion)


$$\Rightarrow \frac{AN}{DN} = \frac{DN}{NB}$$

$$\Rightarrow$$
 DN<sup>2</sup> = AN  $\times$  NB

 $\Rightarrow$  DN<sup>2</sup> = AN  $\times$  DM (As NB = DM)

## Q3:

In the given figure, ABC is a triangle in which  $\angle$  ABC> 90° and AD  $\bot$  CB produced. Prove that AC² = AB² + BC² + 2BC.BD.



### Answer:

Applying Pythagoras theorem in  $\Delta ADB$ , we obtain

$$AB^2 = AD^2 + DB^2 ... (1)$$

Applying Pythagoras theorem in  $\triangle ACD$ , we obtain

$$AC^2 = AD^2 + DC^2$$


$$AC^2 = AD^2 + (DB + BC)^2$$

$$AC^2 = AD^2 + DB^2 + BC^2 + 2DB \times BC$$

$$AC^2 = AB^2 + BC^2 + 2DB \times BC$$
 [Using equation (1)]

## Q4:

In the given figure, ABC is a triangle in which  $\angle$  ABC < 90° and AD  $\bot$  BC. Prove that AC<sup>2</sup> = AB<sup>2</sup> + BC<sup>2</sup> - 2BC.BD.



# Answer:

Applying Pythagoras theorem in  $\triangle ADB$ , we obtain

$$AD^2 + DB^2 = AB^2$$

$$\Rightarrow AD^2 = AB^2 - DB^2 \dots (1)$$



Applying Pythagoras theorem in  $\triangle ADC$ , we obtain

$$AD^2 + DC^2 = AC^2$$

$$AB^2 - BD^2 + DC^2 = AC^2$$
 [Using equation (1)]

$$AB^{2} - BD^{2} + (BC - BD)^{2} = AC^{2}$$

$$AC^2 = AB^2 - BD^2 + BC^2 + BD^2 - 2BC \times BD$$

$$= AB^2 + BC^2 - 2BC \times BD$$


### Q5:

In the given figure, AD is a median of a triangle ABC and AM  $\perp$  BC. Prove that:

$$AC^2 = AD^2 + BC.DM + \left(\frac{BC}{2}\right)^2$$

$$AB^2 = AD^2 - BC.DM + \left(\frac{BC}{2}\right)^2$$
 (ii)

(iii) 
$$AC^2 + AB^2 = 2AD^2 + \frac{1}{2}BC^2$$



#### Answer:

(i) Applying Pythagoras theorem in  $\Delta AMD,$  we obtain

$$AM^2 + MD^2 = AD^2 ... (1)$$

Applying Pythagoras theorem in  $\Delta AMC$ , we obtain

$$AM^2 + MC^2 = AC^2$$

$$AM^2 + (MD + DC)^2 = AC^2$$

$$(AM^2 + MD^2) + DC^2 + 2MD.DC = AC^2$$

$$AD^2 + DC^2 + 2MD.DC = AC^2$$
 [Using equation (1)]

$$DC = \frac{BC}{2}$$

Using the result,



Intelligent sting Intelligent interesting Innovative Learning

Boundary Less Mathematics by PrepOnco
$$AD^{2} + \left(\frac{BC}{2}\right)^{2} + 2MD \cdot \left(\frac{BC}{2}\right) = AC^{2}$$

$$AD^{2} + \left(\frac{BC}{2}\right)^{2} + MD \times BC = AC^{2}$$

(ii) Applying Pythagoras theorem in  $\Delta ABM$ , we obtain

$$AB^2 = AM^2 + MB^2$$

$$= (AD^2 - DM^2) + MB^2$$

$$= (AD^2 - DM^2) + (BD - MD)^2$$

$$= AD^2 - DM^2 + BD^2 + MD^2 - 2BD \times MD$$

$$= AD^2 + BD^2 - 2BD \times MD$$

$$= AD^2 + \left(\frac{BC}{2}\right)^2 - 2\left(\frac{BC}{2}\right) \times MD$$

$$=AD^2 + \left(\frac{BC}{2}\right)^2 - BC \times MD$$

(iii)Applying Pythagoras theorem in ΔABM, we obtain

$$AM^2 + MB^2 = AB^2 ... (1)$$

Applying Pythagoras theorem in  $\Delta$ AMC, we obtain

$$AM^2 + MC^2 = AC^2 ... (2)$$

Adding equations (1) and (2), we obtain

$$2AM^{2} + MB^{2} + MC^{2} = AB^{2} + AC^{2}$$

$$2AM^2 + (BD - DM)^2 + (MD + DC)^2 = AB^2 + AC^2$$

$$2AM^{2}+BD^{2}+DM^{2}-2BD.DM+MD^{2}+DC^{2}+2MD.DC=AB^{2}+AC^{2}$$

$$2AM^{2} + 2MD^{2} + BD^{2} + DC^{2} + 2MD (-BD + DC) = AB^{2} + AC^{2}$$

$$2(AM^2 + MD^2) + (\frac{BC}{2})^2 + (\frac{BC}{2})^2 + 2MD(-\frac{BC}{2} + \frac{BC}{2}) = AB^2 + AC^2$$

$$2AD^2 + \frac{BC^2}{2} = AB^2 + AC^2$$